

333

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 March 2001 (01.03.2001)

PCT

(10) International Publication Number
WO 01/13880 A1

(51) International Patent Classification⁷: A61K 7/48, 7/00

06611 (US). SLAVTCHEFF, Craig, Stephen; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT 06611 (US). ZNAIDEN, Alexander, Paul; Nippon Lever B.V., Utsunomiya Factory/Innovation Centre, 38 Haga-dai, Hago-cho, Haga-gun, Tochigi-ken 321-33 (JP).

(21) International Application Number: PCT/GB00/03031

(22) International Filing Date: 7 August 2000 (07.08.2000)

(25) Filing Language: English

(74) Agent: BAKER, Colin, J.; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).

(26) Publication Language: English

(30) Priority Data:
60/150,481 24 August 1999 (24.08.1999) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for AE, AG, AU, BB, BZ, CA, CY, DM, GB, GD, GH, GM, IE, IL, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW only*): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except AE, AG, AU, BB, BZ, CA, CY, DM, GB, GD, GH, GM, IE, IL, IN, KE, LC, LK, LS, MN, MW, NZ, SD, SG, SL, SZ, TT, TZ, UG, ZA, ZW*): UNILEVER NV [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).

Published:
— With international search report.
— With amended claims.

(71) Applicant (*for IN only*): HINDUSTAN LEVER LIMITED [IN/IN]; Hindustan Lever House, 165/166 Backbay Reclamation, Mumbai 400 020, Maharashtra (IN).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors: GOTT, Robert, Edward; Unilever Home & Personal Care USA, 40 Merritt Boulevard, Trumbull, CT



WO 01/13880 A1

(54) Title: TOWELETTE PRODUCT

(57) Abstract: A disposable towelette product is provided which includes a flexible water-insoluble substrate such as a tissue impregnated with an alpha- or beta- hydroxycarboxylic acid in a cosmetically acceptable carrier vehicle. Impregnated cosmetic compositions in water will have a pH no higher than 6.8. A silicone microemulsion is present to minimize any stickiness resulting from deposition of the hydroxycarboxylic acid by the towelette onto the skin. In the presence of fatty acid group containing surfactants, the silicon microemulsion controls foul odors that the surfactants may emit through hydrolysis at low pH.

TOWELETTE PRODUCT

BACKGROUND OF THE INVENTION

5

Field of the Invention

The present invention relates to single use towelettes for cosmetically delivering alpha-hydroxycarboxylic acids.

10

The Related Art

Alpha-hydroxycarboxylic acids and their derivatives are widely known as providing treatment for the maintenance of a youthful look. These substances are said to control the development of facial fine lines and wrinkles. However, formulation of these substances has been difficult. Among the problems that have been encountered are compatibility with carrier systems, physical stability and skin irritation. In addition, it is particularly difficult to formulate low pH systems.

20

U.S. Patent 5,091,171 (Yu et al.) was one of the first documents to describe the use of alpha-hydroxycarboxylic acids as being effective against the appearance of fine lines and wrinkles. Subsequent thereto a vast amount of literature and many commercial products have been generated based on the efficacy of these materials. Most of the formulations described to date have been of either the cream or lotion type. One problem with these formulations is that they do not always evenly distribute over the applied surfaces. Secondly, any time an active

25

treated surface is followed by cleansing, the active is washed away. Hence, methods of maintaining alpha-hydroxycarboxylic acids on a skin surface are needed which are not susceptible to subsequent cleansing actions. Irritation has also been of great concern because the formulations
5 generally have a low pH.

WO 96/11572 (Moberg) has utilised a variety of acids, including the alpha-hydroxy substance known as lactic acid, in an aqueous hexylene glycol formulation which can be impregnated onto textiles or refreshing
10 napkins. These were employed in order to overcome the problem of microbial growth on skin and also act as a disinfectant.

U.S. Patent 4,828,912 (Hossain et al.) and U.S. Patent 4,764,418 (Kuenn et al.) describe virucidal tissue products directed at controlling disease
15 inducing organisms such as viruses and the common cold. Carboxylic acids such as citric, malic, succinic and benzoic acids are formulated with surfactants and a carrier for impregnation onto facial tissues or other non-woven materials.

20 None of the above disclosures have addressed the issue of stickiness of the alpha-hydroxycarboxylic acids once they have been deposited onto the skin by the towelette. Neither has there been any discussion with respect to odour control which may arise in low pH systems.

25 Accordingly, it is an object of the present invention to provide a product and a method for both cleansing skin and reducing the signs of aging.

Another object of the present invention is to provide a product and a method for depositing alpha-hydroxycarboxylic acids in a manner which avoids the formation of sticky residues on the skin.

- 5 Yet another object of the present invention is to provide a product for delivering alpha-hydroxycarboxylic acids in a formulation which avoids the formation of foul odours.

10 These and other objects of the present invention will become more apparent from the following summary and detailed discussion which follow.

SUMMARY OF THE INVENTION

15

According to the present invention there is provided a towelette product which includes:

- (a) a water-insoluble substrate;
- (b) a cosmetic composition impregnated into the substrate including:
 - 20 (i) an alpha- and beta- hydroxycarboxylic acid;
 - (ii) a silicone microemulsion;
 - (iii) the composition in water having a pH of no higher than about 6.5.

25

Silicone microemulsions of the present invention provide both stability to the composition and counteract any stickiness which may occur when the alpha-hydroxycarboxylic acids are deposited onto the skin.

Surfactants, especially mild surfactants such as those of the amphoteric type may decompose at low pH emitting foul odours. The silicone microemulsions of the present invention have also been found to be useful
5 in counteracting the generation of foul odours.

DETAILED DESCRIPTION OF THE INVENTION

10 The applicants have now surprisingly found that towelettes impregnated with alpha-hydroxycarboxylic acids can be delivered to the skin without imparting any stickiness. This is achieved by the use of silicone microemulsions.

15 Typically creams, lotions and other types of cosmetic vehicles containing the alpha-hydroxycarboxylic acids also contain emollients which are deposited on the skin along with the other actives. These emollients include, for example, esters, hydrocarbons or dimethicone oils. The presence of the emollients obscures any stickiness which may result from
20 evaporation of water or the carrier leaving dried active behind.

Unfortunately, towelette products cannot be impregnated with fluids with too high viscosity as the towelettes will not adequately wet out when the fluid to be impregnated is too thick. However, low viscosity fluids which
25 must be used with towelettes are susceptible to poor emulsion stability and large amounts of emollients are difficult to formulate into these systems. Consequently, actives such as alpha-hydroxycarboxylic acids deposit in a

sticky manner onto skin from the low viscosity fluids spread thereon by the towelettes.

A first necessary aspect of the present invention is that of a substrate. Preferably the substrate is a water insoluble substance. By "water insoluble" is meant that the substrate does not dissolve in or readily break apart upon immersion in water. Another advantage of the substrate in combination with the active is that the former helps the active penetrate. The substrate is also much better than a mere liquid or gel formulation enabling for more accurate application to the skin and the avoidance of sensitive areas, such as the eye where inadvertent application of the astringent composition would cause irritation.

A wide variety of materials may be used as the substrate. The following non-limiting characteristics are desirable: (i) sufficient wet strength for use, (ii) sufficient abrasivity, (iii) sufficient loft and porosity, (iv) sufficient thickness, (v) appropriate size, and (vi) non-reactive with components of the impregnating composition.

Nonlimiting examples of suitable substrates which meet the above criteria include nonwoven substrates, woven substrates, hydroentangled substrates, air entangled substrates and the like. Preferred embodiments employ nonwoven substrates since they are economical and readily available in a variety of materials. By "nonwoven" is meant that the layer is comprised of fibers which are not woven into a fabric but rather are formed into a sheet, particularly a tissue. The fibers can either be random (i.e., randomly aligned) or they can be carded (i.e. combed to be oriented

in primarily one direction). Furthermore, the nonwoven substrate can be composed of a combination of layers of random and carded fibers.

Nonwoven substrates may be comprised of a variety of materials both natural and synthetic. By natural is meant that the materials are derived from plants, animals, insects or byproducts. By synthetic is meant that the materials are obtained primarily from various man-made materials or from material that is usually a fibrous web comprising any of the common synthetic or natural textile-length fibers, or mixtures thereof.

10

Nonlimiting examples of natural materials useful in the present invention are silk fibers, keratin fibers and cellulosic fibers. Nonlimiting examples of keratin fibers include those selected from the group comprising wool fibers, camel hair fibers, and the like. Nonlimiting examples of cellulosic fibers include those selected from the group comprising wood pulp fibers, cotton fibers, hemp fibers, jute fibers, flax fibers, and mixtures thereof. wood pulp fibers are preferred while all cotton fibers (e.g. cotton pads) are normally avoided.

Nonlimiting examples of synthetic materials useful in the present invention include those selected from the group comprising acetate fibers, acrylic fibers, cellulose ester fibers, modacrylic fibers, polyamide fibers, polyester fibers, polyolefin fibers, polyvinyl alcohol fibers, rayon fibers and mixtures thereof. Examples of some of these synthetic materials include acrylics such as Acrilan[®], Creslan[®], and the acrylonitrile-based fiber, Orlon[®]; cellulose ester fibers such as cellulose acetate, Arnel[®] and Acele[®]; polyamides such as Nylons (e.g., Nylon 6, Nylon 66, Nylon 610 and the like); polyesters such as Fortrel[®] Kodel[®] and the polyethylene

25

terephthalate fibers, Dacron® polyolefins such as polypropylene, polyethylene; polyvinyl acetate fibers and mixtures thereof.

Nonwoven substrates made from natural materials consist of webs or
5 sheets most commonly formed on a fine wire screen from a liquid suspension of the fibers.

Substrates made from natural materials useful in the present invention can be obtained from a wide variety of commercial sources. Nonlimiting
10 examples of suitable commercially available paper layers useful herein include Airtex®, an embossed airlaid cellulosic layer having a base weight of about 71 gsy, available from James River Corporation, Green Bay, WI; and Walkisoft®, an embossed airlaid cellulosic having a base weight of about 75 gsy, available from Walkisoft U.S.A., Mount Holly, NC.

15

Nonwoven substrates made from synthetic materials useful in the present invention can also be obtained from a wide variety of commercial sources. Nonlimiting examples of suitable nonwoven layer materials useful herein
20 include HEF 40-047, an apertured hydroentangled material containing about 50% rayon and 50% polyester, and having a basis weight of about 43 grams per square yard (gsy), available from Veratec, Inc., Walpole, MA; HEF 140-102, an apertured hydroentangled material containing about 50% rayon and 50% polyester, and having a basis weight of about 56 gsy, available from Veratec, Inc., Walpole, MA; Novenet® 149-191, a
25 thermo-bonded grid patterned material containing about 69% rayon, about 25% polypropylene, and about 6% cotton, and having a basis weight of about 100 gsy, available from Veratec, Inc., Walpole, MA; HEF Nubtex® 149-801, a nubbed, apertured hydroentangled material, containing about

100% polyester, and having a basis weight of about 70 gsy, available from Veratec, Inc. Walpole, MA; Keybak® 951V, a dry formed apertured material, containing about 75% rayon, about 25% acrylic fibers, and having a basis weight of about 43 gsy, available from Chicopee Corporation, New Brunswick, NJ; Keybak® 1368, an apertured material, containing about 75% rayon, about 5% polyester, and having a basis weight of about 39 gsy, available from Chicopee Corporation, New Brunswick, NJ; Duralace® 1236, an apertured, hydroentangled material, containing about 100% rayon, and having a basis weight from about 40 gsy to about 115 gsy, available from Chicopee Corporation, New Brunswick, NJ; Duralace® 5904, an apertured, hydroentangled material, containing about 100% polyester, and having a basis weight from about 40 gsy to about 115 gsy, available from Chicopee Corporation, New Brunswick, NJ; Sontaro® 8868, a hydroentangled material, containing about 50% cellulose and about 50% polyester, and having a basis weight of about 60 gsy, available from Dupont Chemical Corp.

For the purposes of the present invention the most preferred towelettes are non-woven substrates, especially blends of rayon/polyester in ratios of 10:90 to 90:10, preferably ratios of 20:80 to 80:20, optimally 40:60 to 60:40 by weight. The most preferred towelette is a 70:30 rayon/polyester non-woven wipe article.

The substrate can be made into a wide variety of shapes and forms. Generally the substrate is in single use towelette form. Advantageously, the towelettes are folded in a Z-shaped formation. They may be interleaved with one another but preferably are not interleaved. The Z-fold consists of a center panel flanked by upper and lower wing panels.

The upper and lower wing panels are substantially of equal width and substantially half a width of the center panel. Each towelette is folded medially in a direction orthogonal to that of the Z-shaped formation. Advantageously the size of the towelette may range in length from 10 to 5 40 cm, preferably from 15 to 30 cm, optimally from 18 to 24 cm. The width of the towelette may range from 8 to 30 cm, preferably from 10 to 25 cm, optimally from 15 to 20 cm.


Anywhere from 5 to 100, preferably from 10 to 50 single towelettes may 10 be stored within a dispensing pouch, preferably a moisture impermeable pouch. During storage and between dispensing, the pouch is resealable, usually via an adhesive strip covering a dispensing opening. Single towelette containing pouches may also be employed.

15 The substrates of the present invention may optionally comprise two or more layers, each having a different texture and abrasiveness. The differing textures can result from the use of different combinations of materials or from the use of a substrate having a more abrasive side for exfoliation and a softer, absorbent side for gentle cleansing. In addition, 20 separate layers of the substrate can be manufactured to have different colors, thereby helping the user to further distinguish the surfaces.

A second important element of the present invention is that of an alpha-hydroxycarboxylic acid. By this term is meant not only the acid form but 25 also salts thereof. Typical cationic counterions to form the salt are the alkali metals, alkaline earth metals, ammonium, C2-C8 trialkanolammonium cation and mixtures thereof. The term "alpha-

hydroxycarboxylic acids" include not only hydroxyacids but also alpha-ketoacids and related compounds of polymeric forms of hydroxyacid.

Alpha-hydroxyacids are organic carboxylic acids in which one hydroxyl
5 group is attached to the alpha carbon adjacent the carboxy group. The generic structure is as follows:



10 wherein Ra and Rb are independently an H, F, Cl, Br, alkyl, aralkyl or an aryl group and saturated or unsaturated, isomeric or non-isomeric, straight or branched chain or cyclic form, having 1 to 25 carbon atoms. In addition Ra and Rb may optionally be substituted by OH, CHO, COOH and alkoxy groups having 1 to 9 carbon atoms. The alpha-hydroxyacids
15 may exist as stereoisomers as D, L, and DL forms when Ra and Rb are not identical.

Typical alkyl, aralkyl and aryl groups for Ra and Rb include methyl, ethyl, propyl, isopropyl, butyl, pentyl, octyl, lauryl, stearyl, benzyl and
20 phenyl, etc. The alpha-hydroxyacids of the first group may be subdivided into (1) alkyl alpha-hydroxyacids, (2) aralkyl and aryl alpha-hydroxyacids, (3) polyhydroxy alpha-hydroxyacids, and (4) polycarboxylic alpha-hydroxyacids. The following are representative alpha hydroxyacids in each subgroup:

(1) Alkyl Alpha Hydroxyacids

2-Hydroxyethanoic acid (Glycolic acid, hydroxyacetic acid)

5

2-Hydroxypropanoic acid (Lactic acid)

2-Methyl 2-hydroxypropanoic acid (Methylactic acid)

2-Hydroxybutanoic acid

2-Hydroxypentanoic acid

10

2-Hydroxyhexanoic acid

2-Hydroxyheptanoic acid

2-Hydroxyoctanoic acid

2-Hydroxynonanoic acid

2-Hydroxydecanoic acid

15

2-Hydroxyundecanoic acid

2-Hydroxydodecanoic acid (Alpha hydroxylauric acid)

2-Hydroxytetradecanoic acid (Alpha hydroxymyristic acid)

2-Hydroxyhexadecanoic acid (Alpha hydroxypalmitic acid)

20

2-Hydroxyoctadecanoic acid (Alpha hydroxystearic acid)

2-Hydroxyeicosanoic acid (Alpha hydroxyarachidonic acid)

25

(2) Aralkyl And Aryl Alpha-Hydroxyacids

2-Phenyl 2-hydroxyethanoic acid (Mandelic acid)

2,2-Diphenyl 2-hydroxyethanoic acid (Benzilic acid)
3-Phenyl 2-hydroxypropanoic acid (Phenyllactic acid)
2-Phenyl 2-methyl 2-hydroxyethanoic acid (Atrolactic
acid)

5 2-(4'-Hydroxyphenyl) 2-hydroxyethanoic acid (4-
Hydroxymandelic acid)
2-(4'-Chlorophenyl)-2-hydroxyethanoic acid (4-
Chloromandelic acid)
2-(3'-Hydroxy-4'-methoxyphenyl) 2-hydroxyethanoic
10 acid (3-Hydroxy-4-methoxymandelic acid)
2-(4'-Hydroxy-3'-methoxyphenyl acid)
3-(2'-Hydroxyphenyl) 2-hydroxypropanoic acid
[3(2'-Hydroxyphenyl) lactic acid]
3-(4'-Hydroxyphenyl) 2-hydroxypropanoic acid [3-
15 (4'-Hydroxyphenyl) lactic acid]
2-(3',4'-Dihydroxyphenyl) 2-hydroxyethanoic acid
(3,4-Dihydroxymandelic acid)

(3) Polyhydroxy Alpha-Hydroxyacids

20 2,3-Dihydroxypropanoic acid (Glyceric acid)
2,3,4-Trihydroxybutanoic acid, Isomers; erythronic
acid, threonic acid)
2,3,4,5-Tetrahydroxypentanoic acid (Isomers; ribonic
25 acid, arabinoic acid, xylonic acid, lyxonic acid)
2,3,4,5,6-Pentahydroxyhexanic acid (Isomers; allonic
acid, altronic acid, gluconic acid, mannoic acid,

gulonic acid, idonic acid, galatronic acid, talonic acid)

2,3,4,5,6,7-Hexahydroxyheptanoic acid (Isomers; glucoheptonic acid, galactoheptonic acid etc.)

5

(4) Polycarboxylic Alpha-Hydroxyacids

2-Hydroxypropane-1,3-dioic acid (Tartronic acid)

2-Hydroxybutane,1,4-dioic acid (Malic acid)

10

2,3-Dihydroxybutane-1,4-dioic acid (Tartaric acid)

2-Hydroxy-2-carboxypentane,1,5-dioic acid (Citric acid)

2,3,4,5-Tetrahydroxyhexane,1-5,dioic acid (Isomers: saccharic acid, mucic acid)

15

(5) Lactone Forms

The typical lactone forms are gluconolactone, galactonolactone, glucuronolactone, galacturonolactone, gluconolactone, ribonolactone, saccharic acid lactone, pantoyllactone, glucoheptonolactone, mannonolactone, and galactoheptonolactone.

20

Representative alpha ketoacids useful for the present invention are as follows.

25

2-Ketoethanoic acid (Glyoxylic acid)

Methyl 2-ketoethanoate

2-Ketopropanoic acid (Pyruvic acid)

	Methyl 2-ketopropanoate (Methyl pyruvate)	
	Ethyl 2-ketopropanoate (Ethyl pyruvate)	
	Propyl 2-ketopropanoate (Propyl pyruvate)	
	2-Phenyl-2-ketoethanoic acid (Benzoylformic acid)	
5	Methyl 2-phenyl-2-ketoethanoate (Methyl benzoylformate)	
	Ethyl 2-phenyl-2-ketoethanoate (Ethyl benzoylformate)	
	3-Phenyl-2-ketopropanoic acid (Phenylpyruvic acid)	
10	Methyl 3-phenyl-2-ketopropanoate (Methyl phenylpyruvate)	
	Ethyl 3-phenyl-2-ketopropanoate (Ethyl phenylpyruvate)	
	2-Ketobutanoic acid	
15	2-Ketopentanoic acid	
	2-Ketohexanoic acid	
	2-Ketoheptanoic acid	
	2-Ketooctanoic acid	
	2-Ketododecanoic acid	
20	Methyl 2-ketooctanoate	

II. Dimeric and Polymeric Forms of Hydroxyacids

When two or more molecules of hydroxycarboxylic acids, either identical
25 or non-identical compounds, are reacted chemically to each other, dimeric
or polymeric compounds will be formed. Such dimeric and polymeric
compounds may be classified into three groups, namely (a) acyclic ester,
(b) cyclic ester and (c) miscellaneous dimer and polymer.

Representative acyclic esters of hydroxycarboxylic acids useful for the present invention are those found below.

- 5 Glycolyl glycollate (Glycolic acid glycollate)
 Lactyl lactate (Lactic acid lactate)
 Mandelyl mandellate
 Atrolactyl atrolactate
 Phenyllactyl phenyllactate
10 Benzilyl benzillate
 Glycolyl lactate
 Lactyl glycollate
 Glycolyl glycolyl glycollate
 Lactyl lactyl lactate
15 Lactyl glycolyl lactate
 Glycolyl glycolyl glycolyl glycollate
 Lactyl lactyl lactyl lactate
 Glycolyl lactyl glycolyl lactyl glycollate
 Polyglycolic acid and polylactic acid

20

Amounts of the alpha-hydroxycarboxylic acids may range from about 0.01 to about 20%, preferably from about 0.1 to about 15%, more preferably from about 1 to about 10%, optimally from about 3 to about 8% by weight of the composition which impregnates the substrate.

25

Compositions of the present invention when placed in water will have a pH no higher than about 6.5, preferably from about 6.0 to about 2.0, more preferably from about 5.5 to about 2.5, even more preferably from

about 5.0 to about 3.0, optimally from about 4.5 to about 3.5. The compositions may be placed on the towelettes in a dry state and activated by a consumer through moistening with water. In a preferred embodiment compositions of the present invention are pre-prepared fluid compositions of low viscosity. Typical viscosities may range from 0.5 to 100 cps, preferably from about 2 to about 20 cps at 200C (Brookfield RVT).

The amount of impregnating composition relative to the substrate may range from about 20:1 to 1:20, preferably from 10:1 to about 1:10 and optimally from about 2:1 to about 1:2 by weight.

Typically a humectant is incorporated with compositions of the present invention. Humectants are normally polyols. Representative polyols include glycerin, diglycerin, polyalkylene glycols and more preferably alkylene polyols and their derivatives including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,2-butylene glycol, 1,2,6-hexanetriol, isoprene glycol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. The most preferred is 2-methyl-1,3-propanediol available as MP Diol from the Arco Chemical Company. Amounts of the polyol may range from about 0.5 to about 95%, preferably from about 1 to about 50%, more preferably from about 1.5 to 20%, optimally from about 3 to about 10% by weight of the impregnating composition.

25

An essential further element of product according to the present invention is that of a silicone microemulsion. Average particle size of the silicone material in these microemulsions may range from about 0.01 nm to about

500 nm, preferably from about 1 to about 100 nm, optimally from about 5 to about 50 nm. Particle size may be measured by means of a laser light scattering technique, using a 2600 D Particle Sizer from Malvern Instruments.

5

The microemulsions may be prepared by high shear mechanical mixing of the silicone and water, or by emulsifying the insoluble, non-volatile silicone with water and an emulsifier-mixing the silicone into a heated solution of the emulsifier for instance, or by a combination of mechanical and chemical emulsification.

10

Any surfactant materials either alone or in admixture may be used as emulsifiers in the preparation of the silicone emulsions. Preferred emulsifiers include anionic emulsifiers such as alkylarylsulphonates, e.g. sodium dodecylbenzene sulphonate, alkyl sulphates e.g. sodium lauryl sulphate, alkyl ether sulphates e.g. sodium lauryl ether sulphate nEO, where n is from 1 to 20 alkylphenol ether sulphates e.g. octylphenol ether sulphate nEO where n is from 1 to 20, and sulphosuccinates e.g. sodium dioctylsulphosuccinate.

20

Also suitable are nonionic emulsifiers such as alkylphenol ethoxylates e.g. nonylphenol ethoxylate nEO, where n is from 1 to 50, alcohol ethoxylates e.g. lauryl alcohol nEO, where n is from 1 to 50, ester ethoxylates e.g. polyoxyethylene monostearate where the number of oxyethylene units is from 1 to 30.

25

Silicones which are particularly preferred for purposes of this invention are dimethiconols, these may be linear or branched. Average number

molecular weight may range from about 1,000 to about 1 million, preferably from about 20,000 to about 500,000, optimally from about 40,000 to about 100,000. Microemulsions may be loaded with the silicone at levels ranging from about 1 to about 95%, preferably from about 10 to about 60%, optimally from about 20 to about 40% by weight. Pre-formed microemulsions are available from suppliers such as Dow Corning, General Electric, Union Carbide, Wacker Chemie, Shin Etsu, and Toray Silicone Company. Particularly preferred is a linear dimethiconyl microemulsion at 25% silicone with a maximum particle size of 40 nm, pH 6.5-8 and surfactant combination of dodecylbenzene sulphonic acid triethanolamine/Laureth-24 available from Dow Corning under the trademark DC 2-1870.

Compositions of this invention when in the form of a fluid will usually be provided with a variety of cosmetically acceptable carrier vehicles. Normally the carrier vehicle will be water. Amounts of the carrier vehicle may range from about 0.5 to about 99%, preferably from about 1 to about 80%, more preferably from about 50 to about 70%, optimally from about 65 to 75% by weight of the impregnating composition.

Preservatives can desirably be incorporated into the cosmetic compositions of this invention to protect against the growth of potentially harmful microorganisms. Suitable traditional preservatives for compositions of this invention are alkyl esters of para-hydroxybenzoic acid. Other preservatives which have more recently come into use include hydantoin derivatives, propionate salts, and a variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the preservative

challenge test and to provide product stability. Particularly preferred preservatives are phenoxyethanol, methyl paraben, propyl paraben, imidazolidinyl urea, sodium dehydroacetate and benzyl alcohol. The preservatives should be selected having regard for the use of the composition and possible incompatibilities between the preservatives and other ingredients in the composition. Preservatives are preferably employed in amounts ranging from 0.01% to 2% by weight of the composition.

Compositions of the present invention may further include herbal extracts. Illustrative extracts include Roman Chamomile, Green Tea, Scullcap, Nettle Root, Swertia Japonica, Fennel and Aloe Vera extracts. Amount of each of the extracts may range from about 0.001 to about 1%, preferably from about 0.01 to about 0.5%, optimally from about 0.05 to about 0.2% by weight of a composition.

Minor adjunct ingredients may also be present in the compositions. Among these may be vitamins such as Vitamin E Acetate, Vitamin C, Vitamin A Palmitate, Panthenol and any of the Vitamin B complexes. Anti-irritant agents may also be present including those of steviosides, alpha-bisabolol and glycyhrizzinate salts, each vitamin or anti-irritant agent being present in amounts ranging from about 0.001 to about 1.0%, preferably from about 0.01 to about 0.3% by weight of the composition.

Emulsifiers may also be incorporated into compositions of this invention. These emulsifiers may be anionic, nonionic, cationic, amphoteric and combinations thereof. Useful nonionic type emulsifiers include the C10-C20 fatty alcohol or acid hydrophobes condensed with from 2 to 100

moles of ethylene oxide or propylene oxide per mole of hydrophobe; C2-C10 alkyl phenols condensed with from 2 to 20 moles of alkylene oxide; mono- and di-fatty acid esters of ethylene glycol; fatty acid monoglyceride; sorbitan, mono- and di-C8-C20 fatty acids; block
5 copolymers (ethylene oxide/propylene oxide); and polyoxyethylene sorbitan as well as combinations thereof. Alkyl polyglycosides and saccharide fatty amides (e.g. methyl gluconamides) are also suitable nonionic emulsifiers. Particularly preferred as the emulsifier is a hydrogenated castor wax alkoxylated with 40 moles ethylene oxide,
10 available commercially as Cremophore RH-400®.

Mild emulsifiers of the anionic and amphoteric type may also be employed. Particularly preferred anionic examples include lauroamphoacetate salts and sarcosinate salts. Preferred amphoterics
15 include cocamidopropylbetaine and dimethylbetaine.

Amounts of the emulsifiers may range from about 0.05 to about 20%, preferably from about 0.1 to about 5%, optimally from about 0.5 to about 0.8% by weight.

20

Low pH systems containing fatty acid (C10-C22 alkyl) groups such as lauroamphoacetates have been found to emit foul odours. Although not wishing to be bound by theory, it is believed that hydrolysis of surfactants with fatty acid groups results in cleavage of these groups. Malodorous
25 fatty acids are thereby formed. The applicants have surprisingly found that silicone microemulsions such as dimethiconyl microemulsions effectively inhibit odour formation.

These impregnating compositions of the present invention may involve a range of pH although it is preferred to have a relatively low pH, for instance, a pH from about 2 to about 6.5, preferably from about 2.5 to about 4.5.

5

Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this description indicating amounts of material ought to be understood as modified by the word "about".

- 10 The following examples will more fully illustrate the embodiments of this invention. All parts, percentages and proportions referred to herein and in the appended claims are by weight unless otherwise illustrated.

EXAMPLES 1-8

15

Table I provides a listing of formulations which are suitable for impregnation into a cellulosic substrate forming a towelette. The pH of the resulting composition solutions range from about 2.8 to about 4.0.

EXAMPLE 9

A study was conducted to evaluate the effects of silicones and other substances for stability and reducing stickiness in low pH hydroxycarboxylic acid depositing towelette products. Towelettes were impregnated with a formulation substantially similar to Example 1 except that the dimethiconol microemulsion was replaced with materials reported under Table II at the specified usage level. An amount of 4 grams formulation was impregnated into each towelette (1.8 gram weight; 6 inch by 8 inch size). Stability refers to the phase compatibility of the fluid which impregnated the towelettes. Ratings were 'good' for homogeneous fluids, 'fair, for those with only a small amount of top creaming, and 'poor, for significant phase separation. Stickiness was evaluated by a trained evaluator and measured by sensory finger feel.

TABLE II

Effects of Various Materials on Stability and Stickiness of Low pH Fluids
Delivered by Towelettes

5

Material	CTFA Name	Usage Level	Stability	Effectiveness on stickiness removal (excellent, very good, good, fair, no effect)
Silicones				
Dow Corning 2501	Dimethicone copolyol (wax)	3%	Good	Fair
D/C 2-1788	Dimethicone emulsion	3%	Poor	Good
D/C 2-1310	Dimethicone emulsion	3%	Poor	Good
D/C FB-50	Dimethicone emulsion	3%	Poor	Good
D/C 1664	Dimethicone emulsion	3%	Poor	Good
D/C 2-1870	Dimethiconol microemulsion	3%	Good	Good
D/C 2-8739	Dimethicone microemulsion	3%	Good	Fair
Silwet L-7087	Organosilicone fluid	3%	Good	No effect
Silwax WS-L	Dimethicone copolyol laurate	3%	Good	No effect
Silube CP-I	Dimethicone copolyol phthalate	3%	Good	Fair
Silwax WS	Orfanommodified dimethicone copolyol	3%	Good	Fair
Pecosil PS-100 Q	Dimethicone copolyol phospahte	3%	Good	No effect

D/C antifoam 1510	Dimethicone emulsion	3%	Poor	Good
Others				
Hetester PHA	Propylene glycol isoceteth-3 acetate	3%	Poor	Fair
Lubragel Oil	Glyceryl polymethacralate, propylene glycol, PVM/MA	3%	Good	No effect
Glycerox HE	PEG-7 glyceryl cocoate	3%	Good	No effect
HPS-1180	Polysulfonic acid solution	3%	Good	No effect
		10%	Good	No effect
Lipopeg 4-L	PEG-8 laurate	3%	Good	No effect
Crodesta F-110	Sucrose stearate and sucrose distearate	3%	Fair	No effect
SMEC concentrate	OMC emulsion	3%	Good	No effect
Glycerox 767	PEG 6 capric / caprylic glycerides	3%	Good	No effect
Aqua Keep 10 NF	Sodium polyacrylate (SAP)	1%	Poor	No effect
Arconate HP	Propylene carbonate	3%	Good	No effect
MP diol glycol	2-Methyl 1,3 propanediol	3%	Good	No effect
	2-Methyl 1,3 pentanediol	3%	Good	No effect
Emulgade CM	Cetearyl isonanoate, cetereth-20, cetearyl alcohol, glycerin, cetyl palmitate	3%	Good	No effect

Almost all the non-silicone materials had no beneficial effect on ameliorating stickiness of deposited hydroxycarboxylic acids. Dimethicone emulsions did work well to remove stickiness. However, these emulsions imparted poor stability to the fluid formulations. only
5 dimethiconal microemulsions retained both good physical stability while also effectively combating stickiness.

The foregoing description and examples illustrate selected embodiments of the present invention. In light thereof variations and modifications will be
10 suggested to one skilled in the art, all of which are within the spirit and purview of this invention.

CLAIMS:

1. A towelette product comprising:
 - (a) a water-insoluble substrate;
 - 5 (b) a cosmetic composition impregnated into the substrate including:
 - i) an alpha- and beta- hydroxycarboxylic acid;
 - ii) a silicone microemulsion;
 - 10 iii) the composition in water having a pH no higher than about 6.5

2. A towelette product according to claim 1 wherein the alpha-hydroxycarboxylic acid is selected from the group comprising glycolic,
15 lactic, hydroxyoctanoic acids and mixtures thereof.

3. A towelette product according to claim 1 or claim 2 wherein the pH of the cosmetic composition ranges from about 2.0 to about 6.0.

- 20 4. A towelette product according to claim 1 or claim 2 wherein the pH of the cosmetic composition ranges from about 3.5 to about 4.5.

5. A towelette product according to any of the preceding claims wherein the silicone microemulsion is a dimethiconol microemulsion.
25

6. A towelette product according to Claim 5 wherein the silicone material in the microemulsion has a particle size ranging from about 0.01 to about 500 nm.

7. A towelette product according to claim 5 wherein the silicone material in the microemulsion has a particle size ranging from about 5 to about 50 nm.
- 5 8. A towelette product according any of the preceding claims wherein the hydroxycarboxylic acid is present in an amount from about 0.1 to about 15% by weight of the impregnating vehicle.
9. A towelette product according any of the preceding claims wherein
10 the microemulsion is present in an amount from about 0.1 to about 20% by weight of the impregnating vehicle.
10. A towelette product according to any of the preceding claims wherein the beta-hydroxycarboxylic acid is salicylic acid.
- 15 11. A towelette product comprising:
- (a) a water-insoluble substrate;
 - (b) a composition impregnated onto the substrate
20 comprising:
 - i) an alpha- or beta- hydroxycarboxylic acid;
 - ii) a surfactant containing a C₁₀ - C₂₂ fatty acid group hydrolyzable at low pH; and
 - iii) the composition in water having a pH no higher
25 than 6.8.

AMENDED CLAIMS

[received by the International Bureau on 05 December 2000 (05.12.00);
original claim 11 cancelled ; original claim 1 amended ;
remaining claims unchanged (2 pages)]

1. A towelette product comprising:
 - (a) a water-insoluble substrate;
 - 5 (b) a cosmetic composition impregnated into the substrate including:
 - i) an alpha- or beta- hydroxycarboxylic acid;
 - ii) a silicone microemulsion;
 - 10 iii) the composition in water having a pH no higher than about 6.5

2. A towelette product according to claim 1 wherein the alpha-hydroxycarboxylic acid is selected from the group comprising glycolic, lactic, hydroxyoctanoic acids and mixtures thereof.
- 15 3. A towelette product according to claim 1 or claim 2 wherein the pH of the cosmetic composition ranges from about 2.0 to about 6.0.

4. A towelette product according to claim 1 or claim 2 wherein the pH of the
20 cosmetic composition ranges from about 3.5 to about 4.5.

5. A towelette product according to any of the preceding claims wherein the silicone microemulsion is a dimethiconol microemulsion.

- 25 6. A towelette product according to Claim 5 wherein the silicone material in the microemulsion has a particle size ranging from about 0.01 to about 500 nm.

7. A towelette product according to claim 5 wherein the silicone material in the microemulsion has a particle size ranging from about 5 to about 50 nm.

8. A towelette product according any of the preceding claims wherein the hydroxycarboxylic acid is present in an amount from about 0.1 to about 15% by weight of the impregnating vehicle.
- 5
9. A towelette product according any of the preceding claims wherein the microemulsion is present in an amount from about 0.1 to about 20% by weight of the impregnating vehicle.
- 10 10. A towelette product according to any of the preceding claims wherein the beta-hydroxycarboxylic acid is salicylic acid.

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/GB 00/03031

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K7/48 A61K7/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	DE 199 03 717 A (HENKEL KGAA) 3 August 2000 (2000-08-03) page 3, line 3; claims 1,9	1
A	EP 0 875 233 A (FORT JAMES CORP) 4 November 1998 (1998-11-04) claims 1,30-36	1
A	DE 14 67 886 A (FITZKE, R.) 30 January 1969 (1969-01-30) claims 1,2	1
A	US 3 795 624 A (FEINSTONE W) 5 March 1974 (1974-03-05) claim 1	1
<input type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents :		
A document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search	Date of mailing of the international search report	
22 November 2000	04/12/2000	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016	Authorized officer Beys, E	

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Patent Application No

PCT/GB 00/03031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19903717 A	03-08-2000	AU 4521800 A WO 0044343 A	18-08-2000 03-08-2000
EP 0875233 A	04-11-1998	US 5871763 A	16-02-1999
DE 1467886 A	30-01-1969	NONE	
US 3795624 A	05-03-1974	NONE	